

THE FERMIONIC p -ADIC INTEGRALS ON $\mathbb{Z}_p$ ASSOCIATED WITH EXTENDED q -EULER NUMBERS AND POLYNOMIALS

TAEKYUN KIM

ABSTRACT. The purpose of this paper is to present a systemic study of some families of q -Euler numbers and polynomials of Nörlund's type by using multivariate fermionic p -adic integral on $\mathbb{Z}_p$. Moreover, the study of these higher-order q -Euler numbers and polynomials implies some interesting q -analogue of stirring numbers identities.

§1. Introduction/ Preliminaries

Let p be a fixed odd prime number. Throughout this paper $\mathbb{Z}_p$, $\mathbb{Q}_p$, $\mathbb{C}$ and $\mathbb{C}_p$ will, respectively, denote the ring of p -adic rational integers, the field of p -adic rational numbers, the complex number field and the completion of algebraic closure of $\mathbb{Q}_p$. Let v_p be the normalized exponential valuation of $\mathbb{C}_p$ with $|p|_p = p^{-v_p(p)} = \frac{1}{p}$. When one talks of q -extension, q is variously considered as and indeterminate, a complex number $q \in \mathbb{C}$ or p -adic number $q \in \mathbb{C}_p$. If $q \in \mathbb{C}$, one normally assumes $|q| < 1$. If $q \in \mathbb{C}_p$, one normally assumes $|1 - q|_p < 1$. We use the notation

$$[x]_q = \frac{1 - q^x}{1 - q}, \text{ and } [x]_{-q} = \frac{1 - (-q)^x}{1 + q}, \text{ (see [1-10]).}$$

The q -factorial is defined as $[n]_q! = [n]_q[n-1]_q \cdots [2]_q[1]_q$ and the Gaussian binomial coefficient is also defined by

$$(1) \quad \binom{n}{k}_q = \frac{[n]_q!}{[n-k]_q![k]_q!} = \frac{[n]_q[n-1]_q \cdots [n-k+1]_q}{[k]_q!}, \text{ (see [9]).}$$

Note that

$$\lim_{q \rightarrow 1} \binom{n}{k}_q = \binom{n}{k} = \frac{n!}{(n-k)!k!} = \frac{n(n-1) \cdots (n-k+1)}{k!}.$$

Key words and phrases. : p -adic q -fermionic integral, q -Euler numbers, multivariate integrals.

2000 AMS Subject Classification: 11B68, 11S80

The present Research has been conducted by the research Grant of Kwangwoon University in 2008

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

From (1), we easily note that

$$\binom{n+1}{k}_q = \binom{n}{k-1}_q + q^k \binom{n}{k}_q = q^{n-k} \binom{n}{k-1}_q + \binom{n}{k}_q, \quad (\text{see [5]}).$$

The q -binomial formulae are known that

$$(2) \quad (b; q)_n = (1-b)(1-bq) \cdots (1-bq^{n-1}) = \sum_{i=0}^n \binom{n}{i}_q q^{\binom{i}{2}} (-1)^i b^i,$$

and

$$\frac{1}{(b; q)_n} = \frac{1}{(1-b)(1-bq) \cdots (1-bq^{n-1})} = \sum_{i=0}^{\infty} \binom{n+i-1}{i}_q b^i.$$

We say that f is uniformly differentiable function at a point $a \in \mathbb{Z}_p$, and write $f \in UD(\mathbb{Z}_p)$, if the difference quotient $F_f(x, y) = \frac{f(x)-f(y)}{x-y}$ have a limit $f'(a)$ as $(x, y) \rightarrow (a, a)$. For $f \in UD(\mathbb{Z}_p)$, the fermionic p -adic q -integral on $\mathbb{Z}_p$ is defined as

$$(3) \quad I_{-q}(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-q}(x) = \lim_{N \rightarrow \infty} \frac{1+q}{1+q^{p^N}} \sum_{x=0}^{p^N-1} f(x) (-q)^x, \quad (\text{see [9]}).$$

From (3), we derive the fermionic p -adic integral on $\mathbb{Z}_p$ as follows.

$$(4) \quad \lim_{q \rightarrow 1} I_{-q}(f) = I_{-1}(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-1}(x).$$

For $n \in \mathbb{N}$, let $f_n(x) = f(x+n)$, we have

$$(5) \quad I_{-1}(f_n) = (-1)^n I_{-1}(f) + 2 \sum_{l=0}^{n-1} (-1)^{n-1-l} f(l).$$

From (5), we can easily derive the Euler polynomials, $E_n(x)$, as follows.

$$(6) \quad \int_{\mathbb{Z}_p} e^{(x+y)t} d\mu_{-1} d\mu_{-1}(y) = \frac{2}{e^t + 1} e^{xt} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}, \quad (\text{see [1-23]}).$$

Note that $E_n(0) = E_n$ are called the n -th Euler numbers. Now, we consider the Euler polynomials of Nörlund's type as follows.

$$(7) \quad \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} e^{(x+x_1+\cdots+x_r)t} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r) = \left(\frac{2}{e^t + 1} \right)^r e^{xt} = \sum_{n=0}^{\infty} E_n^{(r)}(x) \frac{t^n}{n!},$$

and

$$(7-1) \quad \left(\frac{e^t + 1}{2} \right)^r e^{xt} = \sum_{n=0}^{\infty} E_n^{(-r)}(x) \frac{t^n}{n!}, \text{ (see [9])}.$$

In the special case $x = 0$, $E_n^{(-r)}(0) = E_n^{(r)}$ and $E_n^{(r)}(0) = E^{(-r)}$ are called the Euler numbers of Nörlund's type. Let $(Eh)(x) = h(x+1)$ be the shift operator. Then the q -difference operator Δ_q is defined as

$$(8) \quad \Delta_q^n = \prod_{i=1}^n (E - q^{i-1}I), \text{ where } (Ih)(x) = h(x), \text{ (see [5, 10])}.$$

From (8), we note that

$$(9) \quad f(x) = \sum_{n \geq 0} \binom{x}{n}_q \Delta_q^n f(0),$$

where

$$\Delta_q^n f(0) = \sum_{k=0}^n \binom{n}{k}_q (-1)^k q^{\binom{k}{2}} f(n-k), \text{ (see [5, 10])}.$$

The q -stirling number of the second kind (as defined by Carlitz) is given by

$$(10) \quad S_2(n, k; q) = \frac{q^{-\binom{k}{2}}}{[k]_q!} \sum_{j=0}^k (-1)^j q^{\binom{j}{2}} \binom{k}{j}_q [k-j]_q^n, \text{ (see [10])}.$$

By (9) and (10), we see that

$$(11) \quad S_2(n, k; q) = \frac{q^{-\binom{k}{2}}}{[k]_q!} \Delta_q^k 0^n, \text{ (see [10])}.$$

In this paper, the q -extension of (7) are variously considered. From these q -extensions, we derive some interesting identities and relations of Euler polynomials and numbers of Nörlund's type. The purpose of this paper is to present a systemic study of some families of q -Euler numbers and polynomials of Nörlund's type by using multivariate fermionic p -adic integral on $\mathbb{Z}_p$

§2. The q -extension of Euler numbers and polynomials of Nörlund type

In this section, we assume that $q \in \mathbb{C}_p$ with $|1 - q|_p < 1$. First, we consider the q -extension of (6) as follows.

$$\begin{aligned} \sum_{n=0}^{\infty} E_{n,q}(x) \frac{t^n}{n!} &= \int_{\mathbb{Z}_p} e^{[x+y]_q t} d\mu_{-q}(x) = \sum_{n=0}^{\infty} \frac{2}{(1-q)^n} \sum_{l=0}^n \left(\frac{\binom{n}{l} (-1)^l q^{lx}}{1 + q^l} \right) \frac{t^n}{n!} \\ &= 2 \sum_{m=0}^{\infty} (-1)^m e^{[m+x]_q t}. \end{aligned}$$

Therefore, we obtain the following lemma.

Lemma 1. For $n \geq 0$, we have

$$(12) \quad E_{n,q}(x) = 2 \sum_{m=0}^{\infty} (-1)^m [m+x]_q^n = \frac{2}{(1-q)^n} \sum_{l=0}^n \frac{\binom{n}{l} (-1)^l q^{lx}}{1+q^l}.$$

From (11), we note that

$$\begin{aligned} [x]_q^n &= \sum_{k=0}^n \binom{n}{k}_q [k]_q! S_2(k, n-k; q) q^{\binom{k}{2}} \\ &= \sum_{k=0}^n [x]_q [x-1]_q \cdots [x-k+1]_q \frac{q^{\binom{k}{2} - \binom{n-k}{2}}}{[n-k]_q!} \Delta_q^{n-k} 0^k \\ &= \sum_{k=0}^n \frac{q^{\binom{k}{2} - \binom{n-k}{2}}}{[n-k]_q!} \Delta_q^{n-k} 0^k \frac{1}{(1-q)^k} \sum_{l=0}^k \binom{k}{l}_q q^{\binom{l}{2}} (-1)^l q^{l(x-k+1)}. \end{aligned}$$

Thus, we have

$$E_{n,q} = \sum_{k=0}^n \frac{q^{\binom{k}{2}} S_2(k, n-k; q)}{(1-q)^k} \sum_{l=0}^k \binom{k}{l}_q q^{\binom{l}{2}} (-1)^l \sum_{m=0}^l \binom{l}{m} (q-1)^m E_{m,q}(1-k).$$

Therefore, we obtain the following theorem.

Theorem 2. For $n \geq 0$, we have

$$E_{n,q} = \sum_{k=0}^n \frac{q^{\binom{k}{2}} S_2(k, n-k; q)}{(1-q)^k} \sum_{l=0}^k \binom{k}{l}_q q^{\binom{l}{2}} (-1)^l \sum_{m=0}^l \binom{l}{m} (q-1)^m E_{m,q}(1-k).$$

Let us consider the q -extension of Eq.(7) as follows.

$$\begin{aligned} E_{n,q}^{(r)}(x) &= \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} [x+x_1+\cdots+x_r]_q^n d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r) \\ (13) \quad &= \frac{2^r}{(1-q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l q^{lx} \left(\frac{1}{1+q^l} \right)^r \\ &= 2^r \sum_{m=0}^{\infty} \binom{m+r-1}{m} (-1)^m [m+x]_q^n. \end{aligned}$$

Let $F_q^{(r)}(t, x) = \sum_{n=0}^{\infty} E_{n,q}^{(r)}(x) \frac{t^n}{n!}$. Then we have

$$(14) \quad F_q^{(r)}(t, x) = 2^r \sum_{m=0}^{\infty} \binom{m+r-1}{m} (-1)^m e^{[m+x]_q t}.$$

In the special case $x = 0$, $E_{n,q}^{(r)}(0) = E_{n,q}^{(r)}$ are called the q -extension of Euler numbers of order r . In the sense of the q -extension of Eq.(7-1), we consider the q -extension of Euler polynomials of Nörlund's type as follows.

$$(15) \quad G_q^{(r)}(t, x) = F_q^{(-r)}(t, x) = \frac{1}{2^r} \sum_{m=0}^r \binom{r}{m} e^{[m+x]_q t} = \sum_{n=0}^{\infty} E_{n,q}^{(-r)}(x) \frac{t^n}{n!}.$$

By (15), we see that

$$E_{n,q}^{(-r)}(x) = \frac{1}{2^r} \sum_{m=0}^r \binom{r}{m} [m+x]_q^n.$$

Therefore, we obtain the following theorem.

Theorem 3. For $r \in \mathbb{N}$, $n \geq 0$, let

$$2^r \sum_{m=0}^{\infty} \binom{m+r-1}{m} (-1)^m e^{[m+x]_q t} = \sum_{n=0}^{\infty} E_{n,q}^{(r)}(x) \frac{t^n}{n!}.$$

Then we have

$$\begin{aligned} E_{n,q}^{(r)}(x) &= \frac{2^r}{(1-q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l q^{lx} \left(\frac{1}{1+q^l} \right)^r \\ &= 2^r \sum_{m=0}^{\infty} \binom{m+r-1}{m} (-1)^m [m+x]_q^n, \end{aligned}$$

and

$$\begin{aligned} E_{n,q}^{(-r)}(x) &= \frac{1}{2^r (1-q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l q^{lx} (1+q^l)^r \\ &= \frac{1}{2^r} \sum_{m=0}^r \binom{r}{m} [m+x]_q^n. \end{aligned}$$

$E_{n,q}^{(-r)}(0) = E_{n,q}^{(-r)}$ are called the q -extension of Euler numbers of Nörlund's type. For $h \in \mathbb{Z}$, $r \in \mathbb{N}$, let us define the extended higher-order q -Euler polynomials as follows.

$$(16) \quad E_{n,q}^{(h,r)}(x) = \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} q^{\sum_{j=1}^r (h-j)x_j} [x + x_1 + \cdots + x_r]_q^n d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r).$$

Then, we have

$$\begin{aligned} E_{n,q}^{(h,r)}(x) &= \frac{2^r}{(1-q)^n} \sum_{l=0}^n \frac{\binom{n}{l} (-1)^l q^{lx}}{(-q^{h-1+l}; q^{-1})_r} = \frac{2^r}{(1-q)^n} \sum_{l=0}^n \frac{\binom{n}{l} (-1)^l q^{lx}}{(-q^{h-r+l}; q)_r} \\ (17) \quad &= 2^r \sum_{m=0}^{\infty} \binom{m+r-1}{m}_q q^{(h-r)m} (-1)^m [x+m]_q^n. \end{aligned}$$

Let $F_q^{(h,r)}(t, x) = \sum_{m=0}^{\infty} E_{n,q}^{(h,r)}(x) \frac{t^n}{n!}$. Then we easily see that

$$(18) \quad F_q^{(h,r)}(t, x) = 2^r \sum_{m=0}^{\infty} q^{(h-r)m} (-1)^m \binom{m+r-1}{m}_q e^{[m+x]_q t}.$$

Therefore we obtain the following theorem.

Theorem 4. For $h \in \mathbb{Z}, n \geq 0$, let

$$2^r \sum_{m=0}^{\infty} q^{(h-r)m} (-1)^m \binom{m+r-1}{m}_q e^{[m+x]_q t} = \sum_{n=0}^{\infty} E_{n,q}^{(h,r)}(x) \frac{t^n}{n!}.$$

Then we have

$$\begin{aligned} E_{n,q}^{(h,r)}(x) &= \frac{2^r}{(1-q)^n} \sum_{l=0}^n \frac{\binom{n}{l} (-1)^l q^{lx}}{(-q^{h-r+l}; q)_r} \\ &= 2^r \sum_{m=0}^{\infty} q^{(h-r)m} (-1)^m \binom{m+r-1}{m}_q [x+m]_q^n. \end{aligned}$$

Now, we define the extended higher-order Nörlund's type q -Euler polynomials as follows.

$$(19) \quad E_{n,q}^{(h,-r)}(x) = \frac{1}{(1-q)^n} \sum_{l=0}^n \frac{\binom{n}{l} (-1)^l q^{lx}}{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} q^{l(x_1+\cdots+x_r)} q^{\sum_{j=1}^r (h-j)x_j} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r)}.$$

From (19), we note that

$$\begin{aligned} (20) \quad E_{n,q}^{(h,-r)}(x) &= \frac{1}{2^r (1-q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l q^{lx} (-q^{h-r+l}; q)_r \\ &= \frac{1}{2^r} \sum_{m=0}^r \binom{r}{m}_q q^{\binom{m}{2}} q^{(h-r)m} [m+x]_q^n. \end{aligned}$$

Let $F_q^{(h,-r)}(t, x) = \sum_{n=0}^{\infty} E_{n,q}^{(h,-r)}(x) \frac{t^n}{n!}$. Then we have

$$(21) \quad F_q^{(h,-r)}(t, x) = \frac{1}{2^r} \sum_{m=0}^r q^{\binom{m}{2}} q^{(h-r)m} \binom{r}{m}_q e^{[m+x]_q t}.$$

Therefore, we obtain the following theorem.

Theorem 5. For $h \in \mathbb{Z}$, $n \geq 0$, $r \in \mathbb{N}$, let

$$\frac{1}{2^r} \sum_{m=0}^{\infty} q^{\binom{m}{2}} q^{(h-r)m} \binom{r}{m}_q e^{[m+x]_q t} = \sum_{n=0}^{\infty} E_{n,q}^{(h,-r)}(x) \frac{t^n}{n!}.$$

Then we have

$$\begin{aligned} E_{n,q}^{(h,-r)}(x) &= \frac{1}{2^r (1-q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l q^{lx} (-q^{h-r+l}; q)_r \\ &= \frac{1}{2^r} \sum_{m=0}^r q^{\binom{m}{2}} q^{(h-r)m} \binom{r}{m}_q [m+x]_q^n. \end{aligned}$$

For $h = r$, we have

$$(22) \quad E_{n,q}^{(r,r)}(x) = \frac{2^r}{(1-q)^n} \sum_{l=0}^n \frac{\binom{n}{l} (-1)^l q^{lx}}{(-q^l; q)_r} = 2^r \sum_{m=0}^{\infty} \binom{m+r-1}{m}_q (-1)^m [x+m]_q^n,$$

and

$$\begin{aligned} (23) \quad E_{n,q}^{(r,-r)}(x) &= \frac{1}{2^r (1-q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l q^{lx} (-q^l; q)_r \\ &= \frac{1}{2^r} \sum_{m=0}^r q^{\binom{m}{2}} \binom{r}{m}_q [m+x]_q^n. \end{aligned}$$

It is easy to see that

$$\begin{aligned} (24) \quad \frac{q^{mx} 2^r}{(-q^{m-r}; q)_r} &= \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} q^{\sum_{j=1}^r (m-j)x_j + mx} d\mu_{-q}(x_1) \cdots d\mu_{-q}(x_r) \\ &= \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} ([x+x_1+\cdots+x_r]_q (q-1) + 1)^m q^{-\sum_{j=1}^r jx_j} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r) \\ &= \sum_{l=0}^m \binom{m}{l} (q-1)^l \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} [x+x_1+\cdots+x_r]_q^l q^{-\sum_{j=1}^r jx_j} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r) \\ &= \sum_{l=0}^m \binom{m}{l} (q-1)^l E_{l,q}^{(0,r)}(x). \end{aligned}$$

By (24), we see that

$$\frac{q^{mx} 2^r}{(-q^{m-r}; q)_r} = \sum_{l=0}^m \binom{m}{l} (q-1)^l E_{l,q}^{(0,r)}(x).$$

It is known that

$$(25) \quad I_{-1}(f_1) + I_{-1}(f) = 2f(0), \text{ where } f_1(x) = f(x+1).$$

From (25), we derive

$$(26) \quad \begin{aligned} & q^{h-1} \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} [x+1+x_1+\cdots+x_r]_q^m q^{\sum_{j=1}^r (h-j)x_j} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r) \\ &= - \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} [x+x_1+\cdots+x_r]_q^n q^{\sum_{j=1}^r (h-j)x_j} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r) \\ &+ 2 \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} [x+x_2+\cdots+x_r]_q^n q^{\sum_{j=1}^{r-1} (h-1-j)x_{j+1}} d\mu_{-1}(x_2) \cdots d\mu_{-1}(x_r). \end{aligned}$$

By (26), we see that

$$(27) \quad q^{(h-1)} E_{n,q}^{(h,r)}(x+1) + E_{n,q}^{(h,r)}(x) = 2E_{n,q}^{(h-1,r-1)}(x).$$

By simple calculation, we see that

$$(28) \quad \begin{aligned} & q^x \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} [x_1+\cdots+x_r+x]_q^n q^{\sum_{j=1}^r (h-j+1)x_j} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r) \\ &= (q-1) \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} [x_1+\cdots+x_r+x]_q^{n+1} q^{\sum_{j=1}^r (h-j)x_j} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r) \\ &+ \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} q^{\sum_{j=1}^r (h-j)x_j} [x_1+\cdots+x_r+x]_q^n d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r). \end{aligned}$$

By (28), we see that

$$q^x E_{n,q}^{(h+1,r)}(x) = (q-1)E_{n+1,q}^{(h,r)}(x) + E_{n,q}^{(h,r)}(x).$$

Therefore, we obtain the following proposition.

Proposition 6. *For $h \in \mathbb{Z}$, $r \in \mathbb{N}$, $n \geq 0$, we have*

$$q^{(h-1)} E_{n,q}^{(h,r)}(x+1) + E_{n,q}^{(h,r)}(x) = 2E_{n,q}^{(h-1,r-1)}(x),$$

and

$$q^x E_{n,q}^{(h+1,r)}(x) = (q-1)E_{n+1,q}^{(h,r)}(x) + E_{n,q}^{(h,r)}(x).$$

Moreover,

$$\frac{q^{mx} 2^r}{(-q^{m-r}; q)_r} = \sum_{l=0}^m \binom{m}{l} (q-1)^l E_{l,q}^{(0,r)}(x).$$

From (22), we note that

$$\begin{aligned}
E_{n,q^{-1}}^{(r,r)}(r-x) &= \frac{2^r}{(1-q^{-1})^n} \sum_{l=0}^n \frac{\binom{n}{l}(-1)^l q^{-l(r-x)}}{(-q^{-l}; q^{-1})_r} \\
&= (-1)^n q^{n+\binom{r}{2}} \frac{2^r}{(1-q)^n} \sum_{l=0}^n \frac{\binom{n}{l}(-1)^l q^{lx}}{(-q^l; q)_r} \\
&= (-1)^n q^{n+\binom{r}{2}} E_{n,q}^{(r,r)}(x).
\end{aligned}$$

Hence, we have

$$\begin{aligned}
&\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} [r-x+x_1+\cdots+x_r]_{q^{-1}}^n q^{-\sum_{j=1}^r (r-j)x_j} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r) \\
&= (-1)^n q^{n+\binom{r}{2}} \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} [x+x_1+\cdots+x_r]_q^n q^{\sum_{j=1}^r (r-j)x_j} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r).
\end{aligned}$$

For $h=r$, we see that

$$E_{n,q^{-1}}^{(r,r)}(0) = (-1)^n q^{n+\binom{r}{2}} E_{n,q}^{(r,r)}(k).$$

From (27), we can also derive

$$q^{r-1} E_{n,q}^{(r,r)}(x+1) + E_{n,q}^{(r,r)}(x) = [2]_q E_{n,q}^{(r-1,r-1)}(x).$$

The stirling numbers of the first kind are defined as

$$(29) \quad \prod_{k=1}^n (1 + [k]_q z) = \sum_{k=0}^n S_1(n, k; q) z^k, \quad (\text{see [10]}),$$

and

$$(30) \quad q^{\binom{m}{2}} \binom{r}{m}_q = \frac{q^{\binom{m}{2}} [r]_q \cdots [r-m+1]_q}{[m]_q!} = \frac{1}{[m]_q!} \prod_{k=0}^{m-1} ([r]_q - [k]_q).$$

It is easy to check that

$$\begin{aligned}
(31) \quad \prod_{k=0}^{n-1} (z - [k]_q) &= z^n \prod_{k=0}^{n-1} \left(1 - \frac{[k]_q}{z}\right) \\
&= \sum_{k=0}^n S_1(n-1, k; q) (-1)^k z^{n-k}.
\end{aligned}$$

By (30) and (31), we see that

$$(32) \quad \prod_{k=0}^{m-1} ([r]_q - [k]_q) = \sum_{k=0}^m S_1(m-1, k; q) (-1)^k [r]_q^{m-k}.$$

By (23) and (32), we obtain the following theorem.

Proposition 7. For $r \in \mathbb{N}$, $n \in \mathbb{Z}_+$, we have

$$E_{n,q}^{(r,-r)}(x) = \frac{1}{2^r [m]_q!} \sum_{m=0}^r \sum_{k=0}^m S_1(m-1, k; q) (-1)^k [r]_q^{m-k} [x+m]_q^n.$$

The generalized Euler numbers and polynomials of Nörlund's type are defined by

$$(33) \quad \frac{2^r}{(e^{w_1 t} + 1)(e^{w_2 t} + 1) \cdots (e^{w_r t} + 1)} = \sum_{n=0}^{\infty} E_n^{(r)}(x|w_1, \dots, w_r) \frac{t^n}{n!},$$

and

$$E_n^{(r)}(w_1, \dots, w_r) = E_n^{(r)}(0|w_1, \dots, w_r).$$

Now, we can also consider the q -extension of (33) as follows. For $w_1, \dots, w_r \in \mathbb{Z}_p$, and $\delta_1, \dots, \delta_r \in \mathbb{Z}$, we define

$$\begin{aligned} & E_{n,q}^{(r)}(x|w_1, \dots, w_r; \delta_1, \dots, \delta_r) \\ &= \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} [xw_1 + \cdots + x_r w_r + x]_q^n d\mu_{-q^{\delta_1}}(x_1) \cdots d\mu_{-q^{\delta_r}}(x_r), \end{aligned}$$

and

$$E_{n,q}^{(r)}(w_1, \dots, w_r; \delta_1, \dots, \delta_r) = E_{n,q}^{(r)}(0|w_1, \dots, w_r; \delta_1, \dots, \delta_r).$$

Thus, we have

$$(34) \quad E_{n,q}^{(r)}(x|w_1, \dots, w_r; \delta_1, \dots, \delta_r) = \sum_{l=0}^n \binom{n}{l} (-1)^l q^{lx} \frac{(1+q^{\delta_1}) \cdots (1+q^{\delta_r})}{(1+q^{\delta_1+lw_1}) \cdots (1+q^{\delta_r+lw_r})}.$$

It seems to be interesting to consider another q -extension of the Nörlund's type generalized Euler numbers and polynomials as follows.

$$(35) \quad \begin{aligned} & E_{n,q}^{*(r)}(x|w_1, \dots, w_r; \delta_1, \dots, \delta_r) \\ &= \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} [w_1 x_1 + \cdots + w_r x_r]_q^n q^{\delta_1 x_1 + \cdots + \delta_r x_r} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r), \end{aligned}$$

and

$$E_{n,q}^{*(r)}(w_1, \dots, w_r; \delta_1, \dots, \delta_r) = E_{n,q}^{*(r)}(0|w_1, \dots, w_r; \delta_1, \dots, \delta_r).$$

From (35), we note that

$$E_{n,q}^{*(r)}(x|w_1, \dots, w_r; \delta_1, \dots, \delta_r) = 2^r \sum_{l=0}^n \frac{\binom{n}{l} (-1)^l q^{lx}}{(1+q^{lw_1+\delta_1}) \cdots (1+q^{lw_r+\delta_r})}.$$

§3. Further Remarks

For $h = 0$, let us consider the polynomial of $E_{n,q}^{(0,r)}(x)$ and $E_{n,q}^{(0,-r)}(x)$ as follows.

$$E_{n,q}^{(0,r)}(x) = \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} [x_1 + \cdots + x_r + x]_q^n q^{-\sum_{j=1}^r jx_j} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r),$$

and

$$E_{n,q}^{(0,-r)} = \sum_{l=0}^n \frac{\binom{n}{l} (-1)^l q^{lx}}{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} q^{l(x_1 + \cdots + x_r)} q^{-\sum_{j=1}^r jx_j} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r)}.$$

Then, we have

$$\begin{aligned} E_{n,q}^{(0,r)}(x) &= \frac{2^r}{(1-q)^n} \sum_{l=0}^n \frac{\binom{n}{l} (-1)^l q^{lx}}{(-q^{l-r}; q)_r} \\ (36) \quad &= 2^r \sum_{m=0}^{\infty} \binom{m+r-1}{m}_q (-1)^m q^{-rm} [m+x]_q^n, \end{aligned}$$

and

$$\begin{aligned} E_{n,q}^{(0,-r)}(x) &= \frac{1}{2^r (1-q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l q^{lx} (-q^{l-r}; q)_r \\ (37) \quad &= \frac{1}{2^r} \sum_{m=0}^r \binom{r}{m}_q q^{\binom{m}{2}} q^{-rm} [m+x]_q^n. \end{aligned}$$

Let $F_q^{(0,r)}(t, x) = \sum_{n=0}^{\infty} E_{n,q}^{(0,r)}(x) \frac{t^n}{n!}$. Then we have

$$F_q^{(0,r)}(t, x) = 2^r \sum_{m=0}^{\infty} \binom{m+r-1}{m}_q (-1)^m q^{-rm} e^{[m+x]_q t},$$

and

$$\begin{aligned} F_q^{(0,-r)}(t, x) &= \sum_{m=0}^{\infty} E_{n,q}^{(0,-r)} \frac{t^n}{n!} \\ &= \frac{1}{2^r} \sum_{m=0}^r \binom{r}{m}_q q^{\binom{m}{2}} q^{-rm} e^{[m+x]_q t}. \end{aligned}$$

Let us consider the following polynomials.

$$\begin{aligned} E_{n,q}^{(h,1)}(x) &= \int_{\mathbb{Z}_p} q^{x_1(h-1)} [x + x_1]_q^n d\mu_{-1}(x_1) \\ (38) \quad &= \frac{2}{(1-q)^n} \sum_{l=0}^n \frac{\binom{n}{l} (-1)^l q^{lx}}{1 + q^{l+h-1}}. \end{aligned}$$

By the simple calculation of fermionic p -adic invariant integral on $\mathbb{Z}_p$, we see that

$$(39) \quad \begin{aligned} & q^x \int_{\mathbb{Z}_p} [x + x_1]_q^n q^{x_1(h-1)} d\mu_{-1}(x_1) \\ &= (q-1) \int_{\mathbb{Z}_p} [x + x_1]_q^{n+1} q^{x_1(h-2)} d\mu_{-1}(x_1) + \int_{\mathbb{Z}_p} [x + x_1]_q^n q^{x_1(h-2)} d\mu_{-1}(x_1). \end{aligned}$$

From (39), we note that

$$q^x E_{n,q}^{(h,1)}(x) = (q-1) E_{n+1,q}^{(h-1,1)}(x) + E_{n,q}^{(h-1,1)}(x).$$

It is not difficult to show that

$$(40) \quad \int_{\mathbb{Z}_p} q^{(h-1)x_1} [x + x_1]_q^n d\mu_{-1}(x_1) = \sum_{j=0}^n \binom{n}{j} [x]_q^{n-j} q^{jx} \int_{\mathbb{Z}_p} [x_1]_q^j q^{(h-1)x_1} d\mu_{-1}(x_1).$$

By (40), we see that

$$E_{n,q}^{(h,1)}(x) = \sum_{j=0}^n \binom{n}{j} [x]_q^{n-j} q^{jx} E_{j,q}^{(h,1)} = \left(q^x E_q^{(h,1)} + [x]_q \right)^n, \quad n \geq 0,$$

where we use the technique method notation by replacing $(E_q^{(h,1)})^n$ by $E_{n,q}^{(h,1)}$, symbolically. From (25), we can also derive

$$(41) \quad \int_{\mathbb{Z}_p} [x + x_1 + 1]_q^n q^{(x_1+1)(h-1)} d\mu_{-1}(x_1) + \int_{\mathbb{Z}_p} [x + x_1]_q^n q^{(h-1)x_1} d\mu_{-1}(x_1) = 2[x]_q^n.$$

Thus, we see that

$$q^{h-1} E_{n,q}^{(h,1)}(x+1) + E_{n,q}^{(h,1)}(x) = 2[x]_q^n.$$

For $x = 0$, we have

$$q^{h-1} \left(q E_q^{(h,1)} + 1 \right)^n + E_{n,q}^{(h,1)} = 2\delta_{n,0},$$

where $\delta_{n,0}$ is the Kronecker symbol.

It is easy to see that

$$E_{0,q}^{(h,1)} = \int_{\mathbb{Z}_p} q^{x_1(h-1)} d\mu_{-1}(x_1) = \frac{2}{1+q^{h-1}} = \frac{2}{[2]_{q^{h-1}}}.$$

From (38), we note that

$$\begin{aligned} E_{n,q^{-1}}^{(h,1)}(1-x) &= \int_{\mathbb{Z}_p} [1-x+x_1]_{q^{-1}}^n q^{-x_1(h-1)} d\mu_{-1}(x_1) \\ &= (-1)^n q^{n+h-1} \frac{2}{(1-q)^n} \sum_{l=0}^n \frac{\binom{n}{l} (-1)^l q^{lx}}{1+q^{l+h-1}} \\ &= (-1)^n q^{n+h-1} E_{n,q}^{(h,1)}(x). \end{aligned}$$

In particular, for $x = 1$, we have

$$E_{n,q^{-1}}^{(h,1)}(0) = (-1)^n q^{n+h-1} E_{n,q}^{(h,1)}(1) = (-1)^{n-1} q^n E_{n,q}^{(h,1)}, \quad \text{for } n \geq 1.$$

REFERENCES

- [1] I. N. Cangul, H. Ozden, Y. Simsek, *A new Approach to q -Genocchi numbers and their interpolations*, Nonlinear Analysis(2008), doi:10.1016/j.na.2008.11.040.
- [2] L. Comtet, *Advanced combinatorics*, Reidel, Dordrecht, 1974.
- [3] E. Deeba, D. Rodriguez, *Stirling's series and Bernoulli numbers*, Amer. Math. Monthly **98** (1991), 423-426.
- [4] M. Cenkci, M. Can and V. Kurt, *p -adic interpolation functions and Kummer-type congruences for q -twisted Euler numbers*, Adv. Stud. Contemp. Math. **9** (2004), 203-216.
- [5] T. Kim, S.-D. Kim, D.-W. Park, *On uniform differentiability and q -Mahler expansions*, Adv. Stud. Contemp. Math. **4** (2001), 35-41.
- [6] T. Kim, *The modified q -Euler numbers and polynomials*, Adv. Stud. Contemp. Math. **16** (2008), 161-170.
- [7] T. Kim, *Euler numbers and polynomials associated with zeta functions*, Abstract and Applied Analysis **2008** (2008), 11 pages(Article ID 581582).
- [8] K. Shiratani, S. Yamamoto, *On a p -adic interpolation function for the Euler numbers and its derivatives*, Mem. Fac. Sci., Kyushu University Ser. A **39** (1985), 113-125.
- [9] T. Kim, *Note on the Euler q -zeta functions*, J. Number Theory(2009), doi:10.1016/j.jnt.2008.10.007.
- [10] T. Kim, *q -Volkenborn integration*, Russ. J. Math. Phys. **9** (2002), 288-299.
- [11] T. Kim, *A Note on p -Adic q -integral on $\mathbb{Z}_p$ Associated with q -Euler Numbers*, Adv. Stud. Contemp. Math. **15** (2007), 133-138.
- [12] T. Kim, *On p -adic interpolating function for q -Euler numbers and its derivatives*, J. Math. Anal. Appl. **339** (2008), 598-608.
- [13] T. Kim, *q -Extension of the Euler formula and trigonometric functions*, Russ. J. Math. Phys. **14** (2007), 275-278.
- [14] T. Kim, *Power series and asymptotic series associated with the q -analog of the two-variable p -adic L -function*, Russ. J. Math. Phys. **12** (2005), 186-196.
- [15] T. Kim, *Non-Archimedean q -integrals associated with multiple Changhee q -Bernoulli polynomials*, Russ. J. Math. Phys. **10** (2003), 91-98.
- [16] T. Kim, *q -Euler numbers and polynomials associated with p -adic q -integrals*, J. Nonlinear Math. Phys. **14** (2007), 15-27.
- [17] H. Ozden, Y. Simsek, S.-H. Rim, I.N. Cangul, *A note on p -adic q -Euler measure*, Adv. Stud. Contemp. Math. **14** (2007), 233-239.
- [18] M. Schork, *Ward's "calculus of sequences", q -calculus and the limit $q \rightarrow -1$* , Adv. Stud. Contemp. Math. **13** (2006), 131-141.
- [19] M. Schork, *Combinatorial aspects of normal ordering and its connection to q -calculus*, Adv. Stud. Contemp. Math. **15** (2007), 49-57.
- [20] Y. Simsek, *On p -adic twisted q - L -functions related to generalized twisted Bernoulli numbers*, Russ. J. Math. Phys. **13** (2006), 340-348.
- [21] Y. Simsek, *Theorems on twisted L -function and twisted Bernoulli numbers*, Advan. Stud. Contemp. Math. **11** (2005), 205-218.
- [22] Y. Simsek, *q -Dedekind type sums related to q -zeta function and basic L -series*, J. Math. Anal. Appl. **318** (2006), 333-351.
- [23] H.J.H. Tuentner, *A Symmetry of power sum polynomials and Bernoulli numbers*, Amer. Math. Monthly **108** (2001), 258-261.

Taekyun Kim

Division of General Education-Mathematics, Kwangwoon University, Seoul 139-701, S. Korea e-mail: tkkim@kw.ac.kr